

11-21-00

A

11/20/00



JC893 U.S. PTO

Docket Number

PI/5-20835B/C1

FILING BY "EXPRESS MAIL" UNDER 37 CFR 1.10

EF080766221US

Express Mail Label Number

November 20, 2000

Date of Deposit

Address to: Assistant Commissioner for Patents  
Box Patent Application  
Washington, DC 20231

JC893 U.S. PTO

09/716656



11/20/00

## UTILITY PATENT APPLICATION TRANSMITTAL AND FEE SHEET

Transmitted herewith for filing under 37 CFR §1.53(b)(1) is a **continuation** of prior Application No. 09/171,700, filed November 20, 2000.

Applicant (or identifier): ROBERT SENN, PETER MAIENFISCH AND PTER WYSS

Title: PESTICIDAL COMPOSITIONS

Enclosed are:

1. ☒ Specification (Including Claims and Abstract) - 35 pages
  2. ☐ Drawings - sheets
  3. ☐ Declaration and Power of Attorney
    - a. ☐ Newly executed (original or copy)
    - b. ☒ Copy from a prior application (signed or with indication that original was signed)
      - i. ☐ Deletion of Inventors  
Signed statement attached deleting inventor(s) named in the prior application
  4. ☒ Incorporation By Reference  
The entire disclosure of the prior application, from which a copy of the Declaration and Power of Attorney is supplied under Box 3b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
  5. ☐ Microfiche Computer Program (appendix)
  6. ☐ Nucleotide and/or Amino Acid Sequence Submission
    - ☐ Computer Readable Copy
    - ☐ Paper Copy
    - ☐ Statement Verifying Identity of Above Copies
  7. ☐ Preliminary Amendment
  8. ☐ Assignment Papers (Cover Sheet & Document(s))
  9. ☐ English Translation of
  10. ☐ Information Disclosure Statement
  11. ☐ Certified Copy of Priority Document(s)
  12. ☒ Return Receipt Postcard
  13. ☐ Other:
- ☒ The right to elect an invention or species that is different from that elected in parent Application No. 09/171,700 in the event of a restriction or election of species requirement that is identical or substantially similar to that made in said parent application is hereby reserved.

09/716656-112000

Filing fee calculation:

- ☐ Before calculating the filing fee, please enter the enclosed Preliminary Amendment.  
☐ Before calculating the filing fee, please cancel claims

Basic Filing Fee								\$	710
Multiple Dependent Claim Fee (\$ 270)								\$	
Foreign Language Surcharge (\$ 130)								\$	
	For	Number Filed		Number Extra		Rate			
Extra Claims	Total Claims	20	-20	2	x	\$	18	=	\$ 36
	Independent Claims	1	-3	0	x	\$	80	=	\$
TOTAL FILING FEE								\$	746


- ☒ Please charge to Credit Card the amount of \$746 as authorized by the enclosed Credit Card Payment Form PTO 2038. An additional copy of this paper is enclosed. The ~~Commissioner is hereby authorized to charge any additional fees under 37 CFR §1.16 and §1.17 which may be required in connection with this application, or credit any overpayment, to the Credit Card.~~

Please address all correspondence to:

William A. Teoli, Jr.  
Syngenta Crop Protection, Inc.  
Patent and Trademark Dept.  
410 Swing Road  
Greensboro, NC 27409

Please direct all telephone calls to the undersigned at the number given below and all telefaxes to (336) 632-2012.

Respectfully submitted,



William A. Teoli, Jr.  
Attorney for Applicants  
Reg. No. 33,104  
Tel. No. (336) 632-7706

Date: November 20, 2000

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IN RE APPLICATION OF  
SENN ET AL.

Art Unit: 1616  
Examiner: A. Robinson

APPLICATION NO: Unassigned  
(Continuation of 09/171,700 filed 11/13/98)

FILED: NOVEMBER 20, 2000

FOR: PESTICIDAL COMPOSITIONS

Assistant Commissioner for Patents  
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Sir:

Prior to substantive examination of this application, kindly amend the claims as follows:

IN THE SPECIFICATION


On page 1, immediately following the title, insert the following;

--This application is a continuation of Serial No. 09/171,700, filed November 13, 1998 (now abandoned).---

IN THE CLAIMS

Cancel, without prejudice, claims 21-22.

1. (Amended) A composition for controlling insects or representatives of the order Acarina, which comprises an effective insecticidal or acaricidal [a] combination [of variable amounts] of one or more compounds of the formula



in which

A is an unsubstituted or [, depending on the possibility of substitution on the ring system,] mono- to tetrasubstituted, aromatic or non-aromatic monocyclic or bicyclic heterocyclic radical, in which the substituents of A are [chosen] selected from the group consisting of C<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>3</sub>alkoxy, halogen, halo-C<sub>1</sub>-C<sub>3</sub>alkyl, cyclopropyl, halocyclopropyl, C<sub>2</sub>-C<sub>3</sub>alkenyl, C<sub>2</sub>-C<sub>3</sub>alkynyl, halo-C<sub>2</sub>-C<sub>3</sub>alkenyl, halo-C<sub>2</sub>-C<sub>3</sub>alkynyl, halo-C<sub>1</sub>-C<sub>3</sub>alkoxy, C<sub>1</sub>-C<sub>3</sub>alkylthio, [Halo-C<sub>1</sub>-C<sub>3</sub>alkylthio] halo-C<sub>1</sub>-C<sub>3</sub>alkylthio, allyloxy, propargyloxy, allylthio, propargylthio, haloallyloxy, haloallylthio, cyano and nitro;

R is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl or C<sub>2</sub>-C<sub>6</sub>alkynyl; and

X is N-NO<sub>2</sub> or N-CN,

in the free form or in salt form, [if appropriate] optionally tautomers thereof, in the free form or salt form, and one or more of the compounds selected from the group consisting of:

- |                            |                          |                          |
|----------------------------|--------------------------|--------------------------|
| (I) aldicarb;              | (XIV) zeta-Cypermethrin; | (XXVII) methamidophos;   |
| (II) azinphos-methyl;      | (XV) deltamethrin;       | (XXVIII) methomyl;       |
| (III) benfuracarb;         | (XVI) diflubenzuron;     | (XXIX) mevinphos;        |
| (IV) bifenthrin;           | (XVII) endosulfan;       | (XXX) parathion;         |
| (V) buprofezin;            | (XVIII) ethiofencarb;    | (XXXI) parathion-methyl; |
| (VI) carbofuran;           | (XIX) fenitrothion;      | (XXXII) phosalone;       |
| (VII) dibutylaminothio;    | (XX) fenobucarb;         | (XXXIII) pirimicarb;     |
| (VIII) cartap;             | (XXI) fenvalerate;       | (XXXIV) propoxur;        |
| (IX) chlorofluazuron;      | (XXII) formothion;       | (XXXV) teflubenzuron;    |
| (X) chloropyrifos;         | (XXIII) methiocarb;      | (XXXVI) terbufos;        |
| (XI) cyfluthrin;           | (XXIV) heptenophos;      | (XXXVII) triazamate;     |
| (XII) lambda-cy-halothrin; | (XXV) imidacloprid;      | (XXXVIII) abamectin;     |
| (XIII) alpha-cypermethrin; | (XXVI) isoprocarb;       | (XXIX) fenobucarb;       |
| (XL) tebufenozide;         | (XLIV) fenpyroximate;    | (XLVIII) pyrimidifen;    |
| (XLI) fipronil;            | (XLV) pyridaben;         | (XLIX) nitenpyram;       |
| (XLII) beta-cyfluthrin;    | (XLVI) fenazaquin;       | (L)[NI-25,] acetamiprid; |
| (XLIII) silafluofen;       | (XLVII) pyriproxyfen;    |                          |

(LI) avermectin B<sub>1</sub> [abamectin]);

(LII) a plant extract which is active against insects;

(LIII) a preparation which comprises nematodes which [are] is active against insects or acarids;

(LIV) a preparation obtainable from *Bacillus subtilis* which is active against insects or acarids;

(LV) a preparation which comprises fungi which [are] is active against insects or acarids;



(CXXXV) omethoate;	(CLVI) sebufos;	(CLXXVII) vamidothion;
(CXXXVI) oxamyl;	(CLVII) sulfotep;	(CLXXVIII) xylylcarb;
(CXXXVII) oxydeme- thon M;	(CLVIII) sulprofos;	(CLXXIX) YI 5301/5302;
(CXXXVIII) oxydeprofos;	(CLIX) tebufenpyrad;	(CLXXX) zetamethrin;
(CXXXIX) permethrin;	(CLX) tebupirimphos;	(CLXXXI) DPX-MP062;
(CXL) phenthoate;	(CLXI) tefluthrin;	(CLXXXII) RH-2485;
(CXLI) phorate;	(CLXII) temephos;	(CLXXXIII) D 2341; or
(CXLII) phosmet;	(CLXIII) terbam;	(CLXXXIV) [XMC (3,5- xylylmethylcarbamate)],
(CXLIII) phoxim;	(CLXIV) tetrachloro- vinphos;	
(CXLIV) pirimiphos M;	(CLXV) thiafenox;	

and at least one formulation auxiliary.

3. (Amended) A composition according to claim 1 [or 2,] in which, in the compound of [the] formula (A), the cyclic base skeleton of A contains 2 to 4 double bonds.

4. (Amended) A composition according to [any one of claims] claim 1 [to 3,] in which, in the compound of the formula (A), the cyclic base skeleton of A contains 1 up to and including 4 heteroatoms.

5. (Amended) A composition according to [any one of claims 1 to] claim 4, in which, in the compound of the formula (A), the cyclic base skeleton of A contains 1, 2 or 3 heteroatoms, [chosen] selected from the group consisting of oxygen, sulfur and nitrogen, not more than one of the heteroatoms contained in the cyclic base skeleton being an oxygen or a sulfur atom.

6. (Amended) A composition according to [any one of claims] claim 1 [to 5,] in which, in the compound of the formula (A) the cyclic base skeleton of A is mono-or disubstituted by substituents [chosen] selected from the group consisting of halogen and C<sub>1</sub>-C<sub>3</sub>alkyl.

7. (Amended) A composition according to [any one of claims 1 to 6,] claim 5 in which, in the compound of the formula (A), the cyclic base skeleton of A is a pyridyl, 1-oxidopyridinio or thiazolyl group.

8. (Amended) A composition according to [any one of claims] claim 1 [to 7,] in which, in the compound of the formula (A), X is N-NO<sub>2</sub>.

10. (Amended) A composition according to [any one of claims] claim 1 [to 9,] which comprises 5-(2-chlorothiazol-5-ylmethyl)-3-methyl-4-nitroimino-perhydro-1,3,5-oxadiazine.

11. (Amended) A composition according to [any one of claims] claim 1 [to 10,] which comprises only one of the compounds (I) to (CLXXXIV).

12. (Amended) A composition according to [any one of claims] claim 1 [to 11,] which comprises pyriproxyfen.

13. (Amended) A composition according to [any one of claims] claim 1 [to 11,] which comprises fipronil.

14. (Amended) A composition according to [any one of claims] claim 1 [to 11,] which comprises endosulfan.

15. (Amended) A composition according to [any one of claims] claim 1 [to 11,] which comprises buprofezin.

16. (Amended) A composition according to [any one of claims] claim 1 [to 11,] which comprises pirimicarb.

17. (Amended) A method of controlling pests, which comprises applying a composition as defined in [any one of claims] claim 1 [to 16,] to the pests or their environment.

18. (Amended) A method [according to claim 17,] for the protection of plant propagation material, which comprises treating the plant propagation material or the site where the propagation material is brought out with the composition according to claim 1.

19. (Amended) A process for the preparation of a composition [comprising at least one auxiliary] as defined in [any one of claims] claim 1 [to 16,] which comprises intimately mixing the active compounds with [the] one or more formulation [auxiliary or] auxiliaries.


REMARKS

Applicant has cancelled claims 21-22, without prejudice, as being in improper "use" format. Applicant reserves the right to file claims to the cancelled subject matter of those claims in applications claiming the benefits hereof. Applicant has amended claims 1, 3-8 and 10-19 to present the claims in more clear and concise terms and to delete all multiple dependencies. No new matter has been added by any of these amendments.

Early and favorable action on the merits is earnestly solicited.

Respectfully submitted,

Syngenta Crop Protection, Inc.  
Patent and Trademark Dept.  
410 Swing Road  
Greensboro, NC 27409  
(336) 632-7706

  
William A. Teoli, Jr.  
Attorney for Applicants  
Reg. No. 33,104

Date: November 20, 2000




Pesticide PESTICIDAL COMPOSITIONS

The present invention relates to a composition which comprises a pesticidal active compound combination, a method for controlling pests with this composition, a process for the preparation of the composition, its use and plant propagation material treated with it, and the use of a compound of the following formula (A) for the preparation of the composition.

Certain mixtures of active compounds are proposed for pest control in the literature. However, the biological properties of these mixtures of known compounds are not completely satisfactory in the field of pest control, and for this reason there is a need to provide further mixtures having synergistic pest control properties, in particular for control of insects and representatives of the order Acarina. This object is achieved according to the invention by providing the present composition.

The invention accordingly relates to a composition for controlling insects and representatives of the order Acarina, which comprises a combination of variable amounts of one or more compounds of the formula



in which

A is an unsubstituted or, depending on the possibility of substitution on the ring system, mono- to tetrasubstituted, aromatic or non-aromatic monocyclic or bicyclic heterocyclic radical, in which the substituents of A are chosen from the group consisting of C<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>3</sub>alkoxy, halogen, halo-C<sub>1</sub>-C<sub>3</sub>alkyl, cyclopropyl, halocyclopropyl, C<sub>2</sub>-C<sub>3</sub>alkenyl, C<sub>2</sub>-C<sub>3</sub>alkynyl, halo-C<sub>2</sub>-C<sub>3</sub>alkenyl, halo-C<sub>2</sub>-C<sub>3</sub>alkynyl, halo-C<sub>1</sub>-C<sub>3</sub>alkoxy, C<sub>1</sub>-C<sub>3</sub>alkylthio, halo-C<sub>1</sub>-C<sub>3</sub>alkylthio, allyloxy, propargyloxy, allylthio, propargylthio, haloallyloxy, haloallylthio, cyano and nitro;

R is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl or C<sub>2</sub>-C<sub>6</sub>alkynyl; and

X is N-NO<sub>2</sub> or N-CN,

in the free form or in salt form, if appropriate tautomers, in the free form or salt form, and one or more of the compounds:

- |  |                          |                           |
|--|--------------------------|---------------------------|
| (I) aldicarb;  | (XVIII) ethiofencarb;    | (XXXV) teflubenzuron;     |
| (II) azinphos-methyl;  | (XIX) fenitrothion;      | (XXXVI) terbufos;         |
| (III) benfuracarb;   | (XX) fenobucarb;         | (XXXVII) triazamate;      |
| (IV) bifenthrin;   | (XXI) fenvalerate;       | (XXXVIII) abamectin;      |
| (V) buprofezin;  | (XXII) formothion;       | (XXXIX) fenobucarb;       |
| (VI) carbofuran;   | (XXIII) methiocarb;      | (XL) tebufenozide;        |
| (VII) dibutylaminothio;  | (XXIV) heptenophos;      | (XLI) fipronil;           |
| (VIII) cartap;   | (XXV) imidacloprid;      | (XLII) beta-cyfluthrin;   |
| (IX) chlorofluazuron;  | (XXVI) isoprocarb;       | (XLIII) silafluofen;      |
| (X) chloropyrifos;   | (XXVII) methamidophos;   | (XLIV) fenpyroximate;     |
| (XI) cyfluthrin;   | (XXVIII) methomyl;       | (XLV) pyridaben;          |
| (XII) lambda-cy-halothrin;   | (XXIX) mevinphos;        | (XLVI) fenazaquin;        |
| (XIII) alpha-cypermethrin;   | (XXX) parathion;         | (XLVII) pyriproxyfen;     |
| (XIV) zeta-Cypermethrin;   | (XXXI) parathion-methyl; | (XLVIII) pyrimidifen;     |
| (XV) deltamethrin;   | (XXXII) phosalone;       | (XLIX) nitenpyram;        |
| (XVI) diflubenzuron;   | (XXXIII) pirimicarb;     | (L) NI-25, acetamiprid;   |
| (XVII) endosulfan;   | (XXXIV) propoxur;        |                           |
| (LI) avermectin B <sub>1</sub> (abamectin);                                      |                          |                           |
| (LII) a plant extract which is active against insects;                           |                          |                           |
| (LIII) a preparation which comprises nematodes which are active against insects; |                          |                           |
| (LIV) a preparation obtainable from <i>Bacillus subtilis</i> ;                   |                          |                           |
| (LV) a preparation which comprises fungi which are active against insects;       |                          |                           |
| (LVI) a preparation which comprises viruses which are active against insects;    |                          |                           |
| (LVII) AC 303 630;   | (LXVII) bendiocarb;      | (LXXVII) carbaryl;        |
| (LVIII) acephate;  | (LXVIII) bensultap;      | (LXXVIII) carbopheno-     |
| (LIX) acrinathrin;   | (LXIX) betacyfluthrin;   | thion;                    |
| (LX) alanycarb;  | (LXX) BPMC;              | (LXXIX) chloethocarb;     |
| (LXI) alphamethrin;  | (LXXI) brofenprox;       | (LXXX) chloroethoxyfos;   |
| (LXII) amitraz;  | (LXXII) bromophos A;     | (LXXXI) chloromephos;     |
| (LXIII) AZ 60541;  | (LXXIII) bufencarb;      | (LXXXII) cis-res-methrin; |
| (LXIV) azinphos A;   | (LXXIV) butocarboxin;    | (LXXXIII) clocythrin;     |
| (LXV) azinphos M;  | (LXXV) butylpyridaben;   | (LXXXIV) clofentezin;     |
| (LXVI) azocyclotin;  | (LXXVI) cadusafos;       | (LXXXV) cyanophos;        |

(LXXXVI) cycloprothrin;	(CXIX) hexaflumuron;	(CL) pyrachlophos;
(LXXXVII) cyhexatin;	(CXX) hexythiazox;	(CLI) pyrada-phenthion;
(LXXXVIII) demeton M;	(CXXI) iprobenfos;	(CLII) pyresmethrin;
(LXXXIX) demeton S;	(CXXII) isofenphos;	(CLIII) pyrethrum;
(XC) demeton-S-methyl;	(CXXIII) isoxathion;	(CLIV) RH 5992;
(XCI) dichlofenthion;	(CXXIV) ivermectin;	(CLV) salithion;
(XCII) dicliphos;	(CXXV) lambda-	(CLVI) sebufos;
(XCIII) diethion;	cyhalothrin;	(CLVII) sulfotep;
(XCIV) dimethoate;	(CXXVI) malathion;	(CLVIII) sulprofos;
(XCV) dimethylvinphos;	(CXXVII) mecarbam;	(CLIX) tebufenpyrad;
(XCVI) dioxathion;	(CXXVIII) mesulfenphos;	(CLX) tebupirimphos;
(XCVII) edifenphos;	(CXXIX) metaldehyde;	(CLXI) tefluthrin;
(XCVIII) emamectin;	(CXXX) metolcarb;	(CLXII) temephos;
(XCIX) esfenvalerate;	(CXXXI) milbemectin;	(CLXIII) terbam;
(C) ethion;	(CXXXII) moxidectin;	(CLXIV) tetrachloro-
(CI) ethofenprox;	(CXXXIII) naled;	vinphos;
(CII) ethoprophos;	(CXXXIV) NC 184;	(CLXV) thiafenox;
(CIII) etrimphos;	(CXXXV) omethoate;	(CLXVI) thiodicarb;
(CIV) fenamiphos;	(CXXXVI) oxamyl;	(CLXVII) thiofanox;
(CV) fenbutatin oxide;	(CXXXVII) oxydeme-	(CLXVIII) thionazin;
(CVI) fenothiocarb;	thon M;	(CLXIX) thuringiensin;
(CVII) fenpropathrin;	(CXXXVIII) oxydeprofos;	(CLXX) tralomethrin;
(CVIII) fenpyrad;	(CXXXIX) permethrin;	(CLXXI) triarthen;
(CIX) fenthion;	(CXL) phenthoate;	(CLXXII) triazophos;
(CX) fluazinam;	(CXLI) phorate;	(CLXXIII) triazuron;
(CXI) flucycloxuron;	(CXLII) phosmet;	(CLXXIV) trichlorofon;
(CXII) flucythrinate;	(CXLIII) phoxim;	(CLXXV) triflumuron;
(CXIII) flufenoxuron;	(CXLIV) pirimiphos M;	(CLXXVI) trimethacarb;
(CXIV) flufenprox;	(CXLV) pirimiphos A;	(CLXXVII) vamidothion;
(CXV) fonophos;	(CXLVI) promecarb;	(CLXXVIII) xylylcarb;
(CXVI) fosthiazate;	(CXLVII) propaphos;	(CLXXIX) YI 5301/5302;
(CXVII) fubfenprox;	(CXLVIII) prothiofos;	(CLXXX) zetamethrin;
(CXVIII) HCH;	(CXLIX) prothoate;	(CLXXXI) DPX-MP062;

0971656-11000

(CLXXXII) RH-2485; (CLXXXIII) D 2341; or (CLXXXIV) XMC (3,5-xylyl-methylcarbamate),  
and at least one auxiliary.

Compounds of the formula (A) are described in EP-A-580553.

The compounds (I) to (CLXXXIV) are known to the expert. In particular

- (I) 2-methyl-2-(methylthio)propionaldehyde O-methylcarbamoyloxime (aldicarb) is known from The Pesticide Manual, 9<sup>th</sup> Ed. (1991), The British Crop Protection Council, London, page 16;
- (II) S-3,4-dihydro-4-oxo-1,2,3-benzotriazin-3-ylmethyl O,O-dimethyl phosphorodithioate (azinphos-methyl) is known from The Pesticide Manual, 9<sup>th</sup> Ed. (1991), The British Crop Protection Council, London, page 46;
- (III) ethyl N-[2,3-dihydro-2,2-dimethylbenzofuran-7-yloxycarbonyl(methyl)aminothio]-N-isopropyl- $\beta$ -alaninate (benfuracarb) is known from The Pesticide Manual, 9<sup>th</sup> Ed. (1991), The British Crop Protection Council, London, page 57;
- (IV) 2-methylbiphenyl-3-ylmethyl (Z)-(1RS)-cis-3-(2-chloro-3,3,3-trifluoroprop-1-enyl)-2,2-dimethylcyclopropanecarboxylate (bifenthrin) is known from The Pesticide Manual, 9<sup>th</sup> Ed. (1991), The British Crop Protection Council, London, page 73;
- (V) 2-tert-butylimino-3-isopropyl-5-phenyl-1,3,5-thiadiazian-4-one (buprofezin) is known from The Pesticide Manual, 9<sup>th</sup> Ed. (1991), The British Crop Protection Council, London, page 105;
- (VI) 2,3-dihydro-2,2-dimethylbenzofuran-7-yl methylcarbamate (carbofuran) is known from The Pesticide Manual, 9<sup>th</sup> Ed. (1991), The British Crop Protection Council, London, page 126;
- (VII) 2,3-dihydro-2,2-dimethylbenzofuran-7-yl (dibutylaminothio)-methylcarbamate (carbosulfan) is known from The Pesticide Manual, 9<sup>th</sup> Ed. (1991), The British Crop Protection Council, London, page 129;
- (VIII) S,S'-(2-dimethylaminotrimethylene) bis(thiocarbamate) (cartap) is known from The Pesticide Manual, 9<sup>th</sup> Ed. (1991), The British Crop Protection Council, London, page 132;
- (IX) 1-[3,5-dichloro-4-(3-chloro-5-trifluoromethyl-2-pyridyloxy)phenyl]-3-(2,6-difluorobenzoyl)-urea (chlorfluazuron) is known from The Pesticide Manual, 9<sup>th</sup> Ed. (1991), The British Crop Protection Council, London, page 143;

- (X) O,O-diethyl O-3,5,6-trichloro-2-pyridyl phosphorothioate (chlorpyrifos) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 166;
- (XI) (RS)- $\alpha$ -cyano-4-fluoro-3-phenoxybenzyl (1RS)-cis-trans-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (cyfluthrin) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 198;
- (XII) Mixture of (S)- $\alpha$ -cyano-3-phenoxybenzyl Z-(1R)-cis-3-(2-chloro-3,3,3-trifluoropropenyl)-2,2-dimethylcyclopropanecarboxylate and (R)- $\alpha$ -cyano-3-phenoxybenzyl Z-(1S)-cis-3-(2-chloro-3,3,3-trifluoropropenyl)-2,2-dimethylcyclopropanecarboxylate (lambda-cyhalothrin) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 203;
- (XIII) Racemate of (S)- $\alpha$ -cyano-3-phenoxybenzyl (1R)-cis-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate and (R)- $\alpha$ -cyano-3-phenoxybenzyl (1S)-cis-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (alpha-cypermethrin) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 210;
- (XIV) A mixture of (S)- $\alpha$ -cyano-3-phenoxybenzyl (1RS)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate and (S)- $\alpha$ -cyano-3-phenoxybenzyl (3RS)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (zeta-cypermethrin) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, page 265;
- (XV) (S)- $\alpha$ -cyano-3-phenoxybenzyl (1R)-cis-3-(2,2-dibromovinyl)-2,2-dimethylcyclopropanecarboxylate (deltamethrin) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 232;
- (XVI) (4-chlorophenyl)-3-(2,6-difluorobenzoyl)urea (diflubenzuron) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 281;
- (XVII) (1,4,5,6,7,7-hexachloro-8,9,10-trinorborn-5-en-2,3-ylenebismethylene) sulfite (endosulfan) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 332;
- (XVIII)  $\alpha$ -ethylthio-o-tolyl methylcarbamate (ethiofencarb) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 343;

- (XIX) O,O-dimethyl O-4-nitro-m-tolyl phosphorothioate (fenitrothion) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 369;
- (XX) 2-sec-butylphenyl methylcarbamate (fenobucarb) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 371;
- (XXI) (RS)- $\alpha$ -cyano-3-phenoxybenzyl (RS)-2-(4-chlorophenyl)-3-methylbutyrate (fenvalerate) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 388;
- (XXII) [formyl(methyl)carbamoylmethyl] O,O-dimethyl phosphorodithioate (formothion) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 440;
- (XXIII) 4-methylthio-3,5-xylyl methylcarbamate (methiocarb) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, page 677;
- (XXIV) 7-chlorobicyclo[3.2.0]hepta-2,6-dien-6-yl dimethylphosphate (heptenophos) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, page 467;
- (XXV) (6-chloro-3-pyridylmethyl)-N-nitroimidazolidin-2-ylidenamine (imidacloprid) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 491;
- (XXVI) 2-isopropylphenyl methylcarbamate (isoprocab) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 504;
- (XXVII) O,S-dimethyl phosphoramidothioate (methamidophos) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 563;
- (XXVIII) S-methyl N-(methylcarbamoyloxy)thioacetimidate (methomyl) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 570;
- (XXIX) methyl 3-(dimethoxyphosphinoyloxy)but-2-enoate (mevinphos) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 592;
- (XXX) O,O-diethyl O-4-nitrophenyl phosphorothioate (parathion) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 648;
- (XXXI) O,O-dimethyl O-4-nitrophenyl phosphorothioate (parathion-methyl) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 650;

09716556 112000

- (XXXII) S-6-chloro-2,3-dihydro-2-oxo-1,3-benzoxazol-3-ylmethyl O,O-diethyl phosphordithioate (phosalone) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 674;
- (XXXIII) 2-dimethylamino-5,6-dimethylpyrimidin-4-yl dimethylcarbamate (pirimicarb) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 690;
- (XXXIV) 2-isopropoxyphenyl methylcarbamate (propoxur) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 727;
- (XXXV) 1-(3,5-dichloro-2,4-difluorophenyl)-3-(2,6-difluorobenzoyl)urea (teflubenzuron) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 790;
- (XXXVI) S-tert-butylthiomethyl O,O-dimethyl phosphorodithioate (terbufos) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 795;
- (XXXVII) ethyl (3-tert.-butyl-1-dimethylcarbamoyl-1H-1,2,4-triazol-5-yl-thio)acetate (triazamate) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 1006;
- (XXXVIII) abamectin is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 3;
- (XXXIX) 2-sec-butylphenyl methylcarbamate (fenobucarb) is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 371;
- (XL) N-tert-butyl-N'-(4-ethylbenzoyl)-3,5-dimethylbenzohydrazide (tebufenozide) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 943;
- (XLI) ( $\pm$ )-5-amino-1-(2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl)-4-trifluoromethyl-sulfinylpyrazole-3-carbonitril (fipronil) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 463;
- (XLII)  $\alpha$ -cyano-4-fluoro-3-phenoxybenzyl 3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropane-carboxylate (beta-cyfluthrin) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 250;
- (XLIII) (4-ethoxyphenyl)[3-(4-fluoro-3-phenoxyphenyl)propyl](dimethyl)silane (silaflluofen) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 912;

- (XLIV) tert-butyl (E)- $\alpha$ -(1,3-dimethyl-5-phenoxy-pyrazol-4-yl-methylenamino-oxy)-p-toluate (fenpyroximate) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 450;
- (XLV) 2-tert-butyl-5-(4-tert.-butylbenzylthio)-4-chloropyridazin-3(2H)-one (pyridaben) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 879;
- (XLVI) 4-tert-butylphenethyl quinazolin-4-yl ether (fenazaquin) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 426;
- (XLVII) 4-phenoxyphenyl (RS)-2-(pyridyloxy)propyl ether (pyriproxyfen) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 887;
- (XLVIII) 5-chloro-N-{2-[4-(2-ethoxyethyl)-2,3-dimethylphenoxy]ethyl}-6-ethylpyrimidin-4-amine (pyrimidifen) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 887;
- (XLIX) (E)-N-(6-chloro-3-pyridylmethyl)-N-ethyl-N'-methyl-2-nitrovinylidenediamine (nitenpyram) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 736;
- (L) (E)-N<sup>1</sup>-[(6-chloro-3-pyridyl)methyl]-N<sup>2</sup>-cyano-N<sup>1</sup>-methylacetamidine (NI-25, acetamiprid) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 730;
- (LI) avermectin B<sub>1</sub> (abamectin) is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 1;
- (LII) a plant extract which is active against insects, in particular (2R,6aS,12aS)-1,2,6,6a,12,12a-hexhydro-2-isopropenyl-8,9-dimethoxy-chromeno[3,4-b]furo[2,3-h]-chromen-6-one (rotenone), is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 758; and an extract from Azadirachta indica, in particular azadirachtin, is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 54; and
- (LIII) a preparation which comprises nematodes which are active against insects, preferably Heterorhabditis bacteriophora, Heterorhabditis megidis is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 560; one comprising Steinernema feltiae is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 920, and one comprising Steinernema



scapterisci is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 921;

(LIV) a preparation obtainable from *Bacillus subtilis* is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 62; or one obtainable from a *Bacillus thuringiensis* strain, with the exception of compounds isolated from GC91 or from NCTC11821, is known;

(LV) a preparation which comprises fungi which are active against insects, preferably *Verticillium lecanii*, is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 1040; one comprising *Beauveria brogniartii* is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 69; and one comprising *Beauveria bassiana* is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 69;

(LVI) a preparation which comprises viruses which are active against insects, preferably *Neodiprion sertifer* NPV, is known from The Pesticide Manual, 9<sup>th</sup>Ed. (1991), The British Crop Protection Council, London, Page 616; one comprising *Mamestra brassicae* NPV is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 633; and one comprising *Cydia pomonella* granulosus virus is known from The Pesticide Manual, 10<sup>th</sup>Ed. (1994), The British Crop Protection Council, London, Page 247;

(CLXXXI) indeno[1,2-e] [1,3,4] oxadiazine-4a (3H)-carboxylic acid, 7-chloro-2,5-dihydro-2-[[[(methoxycarbonyl)[(4-trifluoromethoxy)phenyl]amino]carbonyl]-, methyl ester (DPX-MP062), is known from the Brighton Crop Protection Conference, 1996, 449- 456;

(CLXXXII) N'-tert-butyl-N'-(3,5-dimethylbenzoyl)-3-methoxy-2-methylbenzohydrazide (RH-2485, methoxyfenozide) is known from the Brighton Crop Protection Conference, 1996, 481- 486; and

(CLXXXIII) N'-[4-methoxybiphenyl-3-yl]hydrazinecarboxylic acid isopropyl ester (D 2341) is known from the Brighton Crop Protection Conference, 1996, 487- 493.

The compounds of the formula (A) can in some cases be in the form of tautomers. For example, if R is hydrogen, corresponding compounds of the formula (A), i.e. those having a 3-H-4-imino-perhydro-1,3,5-oxadiazine partial structure, can be in equilibrium with the particular tautomers which contain a 4-amino-1,2,5,6-tetrahydro-1,3,5-oxadiazine partial structure. Compounds of the formula (A) above and below are accordingly also to be un-

derstood as meaning, where appropriate, corresponding tautomers, even if the latter are not mentioned specifically in each case.

Compounds of the formula (A) which contain at least one basic centre can form, for example, acid addition salts. These are formed, for example, with strong inorganic acids such as mineral acids, for example perchloric acid, sulfuric acid, nitric acid, nitrous acid, a phosphoric acid or a hydrogen halide acid, with strong organic carboxylic acids, such as C<sub>1</sub>-C<sub>4</sub>alkanecarboxylic acids which are unsubstituted or substituted, for example by halogen, for example acetic acid, such as dicarboxylic acids which are saturated or unsaturated, for example oxalic, malonic, succinic, maleic, fumaric or phthalic acid, such as hydroxycarboxylic acids, for example ascorbic, lactic, malic, tartaric or citric acid, or such as benzoic acid, or with organic sulfonic acids, such as C<sub>1</sub>-C<sub>4</sub>alkane- or arylsulfonic acids which are unsubstituted or substituted, for example by halogen, for example methane- or p-toluenesulfonic acid. Compounds of the formula (A) with at least one acid group can furthermore form salts with bases. Suitable salts with bases are, for example, metal salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, piperidine, pyrrolidine, a mono-, di- or tri-lower alkylamine, for example ethyl-, diethyl-, triethyl- or dimethyl-propyl-amine, or a mono- di- or trihydroxy-lower alkylamine, for example mono-, di- or triethanolamine. If appropriate, corresponding inner salts can furthermore be formed. Agrochemically advantageous salts are preferred in the context of the invention. As a result of the close relationship between the compounds of the formula (A) in the free form and in the form of their salts, the free compounds of the formula (A) and their salts above and below are also appropriately and expediently to be understood as meaning, where appropriate, the corresponding salts and the free compounds of the formula (A). The same applies correspondingly to tautomers of compounds of the formula (A) and salts thereof. In each case the free form is in general preferred.

Compositions which comprise the compound of the formula (A) in the free form are preferred in the context of the present invention.

The general terms used above and below have the meanings given below, unless defined otherwise.

Heteroatoms in the cyclic base skeleton of the heterocyclic radical R are all elements of the Periodic Table which can form at least two covalent bonds. N and S are preferred.

Halogen - as a group per se and as a structural element of other groups and compounds, such as of haloalkyl, haloalkylthio, haloalkoxy, halocyclopropyl, haloalkenyl, haloalkynyl, haloallyloxy and haloallylthio - is fluorine, chlorine, bromine or iodine, in particular fluorine, chlorine or bromine, especially fluorine or chlorine, in particular chlorine..

Unless defined otherwise, carbon-containing groups and compounds in each case contain 1 up to and including 6, preferably 1 up to and including 3, in particular 1 or 2, carbon atoms.

Cycloalkyl is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, preferably cyclopropyl.

Alkyl - as a group per se and as a structural element of other groups and compounds, such as of phenylalkyl, haloalkyl, alkoxy, haloalkoxy, alkylthio and haloalkylthio - is, in each case taking into due consideration the number of carbon atoms, defined from case to case, contained in the corresponding group or compound, either straight-chain, i.e. methyl, ethyl, propyl, butyl, pentyl or hexyl, or branched, for example, isopropyl, isobutyl, sec-butyl, tert-butyl, isopentyl, neopentyl or isohexyl.

Alkenyl, haloalkenyl, alkynyl and haloalkynyl are straight-chain or branched and in each case contain two or preferably one unsaturated carbon-carbon bond(s). The double or triple bonds of this substituent are preferably separated from the remaining part of the compound of the formula (A) by at least one saturated carbon atom. Examples are allyl, methallyl, but-2-enyl, but-3-enyl, propargyl, but-2-ynyl and but-3-ynyl.

Halogen-substituted carbon-containing groups and compounds, such as haloalkyl, haloalkylthio, haloalkoxy, halocyclopropyl, haloalkenyl, haloalkynyl, haloallyloxy and haloallylthio, can be partly halogenated or perhalogenated, and in the case of multiple halogenation the halogen substituents can be identical or different. Examples of haloalkyl - as a group per se and as a structural element of other groups and compounds, such as of haloalkylthio and haloalkoxy - are methyl which is mono- to trisubstituted by fluorine, chlorine and/or bromine, such as  $\text{CHF}_2$  or  $\text{CF}_3$ ; ethyl which is mono- to pentasubstituted by fluorine, chlorine and/or bromine, such as  $\text{CH}_2\text{CF}_3$ ,  $\text{CF}_2\text{CF}_3$ ,  $\text{CF}_2\text{CCl}_3$ ,  $\text{CF}_2\text{CHCl}_2$ ,  $\text{CF}_2\text{CHF}_2$ ,  $\text{CF}_2\text{CFCl}_2$ ,  $\text{CF}_2\text{CHBr}_2$ ,  $\text{CF}_2\text{CHClF}$ ,  $\text{CF}_2\text{CHBrF}$  or  $\text{CClFCHClF}$ ; propyl or isopropyl which are mono- to heptasubsti-

tuted by fluorine, chlorine and/or bromine, such as  $\text{CH}_2\text{CHBrCH}_2\text{Br}$ ,  $\text{CF}_2\text{CHFCH}_2\text{F}$ ,  $\text{CH}_2\text{CF}_2\text{CF}_3$ ,  $\text{CF}_2\text{CF}_2\text{CF}_3$  or  $\text{CH}(\text{CF}_3)_2$ ; and butyl or one of its isomers which are mono- to nonasubstituted by fluorine, chlorine and/or bromine, such as  $\text{CF}(\text{CF}_3)\text{CHFCH}_2\text{F}$ ,  $\text{CF}_2(\text{CF}_2)_2\text{CF}_3$  or  $\text{CH}_2(\text{CF}_2)_2\text{CF}_3$ . Examples of haloalkenyl are 2,2-difluoroethen-1-yl, 2,2-dichloroethen-1-yl, 2-chloroprop-1-en-3-yl, 2,3-dichloroprop-1-en-3-yl and 2,3-dibromoprop-1-en-3-yl. Examples of haloalkynyl are 2-chloroprop-1-yn-3-yl, 2,3-dichloroprop-1-yn-3-yl and 2,3-dibromoprop-1-yn-3-yl. Examples of halocyclopropyl are 2-chlorocyclopropyl, 2,2-difluorocyclopropyl and 2-chloro-2-fluoro-cyclopropyl. Examples of haloallyloxy are 2-chloroprop-1-en-3-yloxy, 2,3-dichloroprop-1-en-3-yloxy and 2,3-dibromoprop-1-en-3-yloxy. Examples of haloallylthio are 2-chloroprop-1-en-3-ylthio, 2,3-dichloroprop-1-en-3-ylthio and 2,3-dibromoprop-1-en-3-ylthio.

In phenylalkyl, an alkyl group bonded to the remaining part of the compound of the formula (A) is substituted by a phenyl group, the alkyl group preferably being straight-chain and the phenyl group preferably being bonded to the alkyl group in a position higher than the  $\alpha$ -position, in particular in the  $\omega$ -position; examples are benzyl, 2-phenylethyl and 4-phenylbutyl.

Particularly preferred compounds of the formula (A) are those in which

- (1) R is hydrogen,  $\text{C}_1$ - $\text{C}_4$ alkyl,  $\text{C}_3$ - $\text{C}_6$ cycloalkyl,  $\text{C}_2$ - $\text{C}_6$ alkenyl or  $\text{C}_2$ - $\text{C}_6$ alkynyl, in particular H or  $\text{CH}_3$ ;
- (2) the cyclic base skeleton of A contains 2 to 4, preferably conjugated, double bonds, preferably contains 2, preferably conjugated double bonds, and in particular has aromatic character,
- (3) the cyclic base skeleton of A contains 1 up to and including 4, in particular 1 up to and including 3, especially 1 or 2, heteroatoms, particularly preferably 2 heteroatoms;
- (4) the cyclic base skeleton of A contains 1, 2 or 3 heteroatoms chosen from the group consisting of oxygen, sulfur and nitrogen, not more than one of the heteroatoms contained in the cyclic base skeleton being an oxygen or sulfur atom, preferably at least one nitrogen atom;

(5) a compound of the formula (A) , in which A is mono- or disubstituted by substituents chosen from the group consisting of halogen and C<sub>1</sub>-C<sub>3</sub>alkyl, in particular by halogen, especially by chlorine;

(6) a compound of the formula (A) in which the cyclic base skeleton of A is a pyridyl, 1-oxidopyridinio or thiazolyl group,  
preferably the cyclic base skeleton of A is a pyrid-3-yl-, 1-oxido-3-pyridinio or thiazol-5-yl group,

in particular A is a pyrid-3-yl, 2-halopyrid-5-yl, 2,3-dihalopyrid-5-yl, 2-C<sub>1</sub>-C<sub>3</sub>alkylpyrid-5-yl, 1-oxido-3-pyridinio, 2-halo-1-oxido-5-pyridinio, 2,3-dihalo-1-oxido-5-pyridinio or 2-halothiazol-5-yl group,

A is especially a pyrid-3-yl, 2-halopyrid-5-yl, 2-halo-1-oxido-5-pyridinio or 2-halothiazol-5-yl group,

preferably A is a 2-chloropyrid-5-yl, 2-methylpyrid-5-yl, 1-oxido-3-pyridinio, 2-chloro-1-oxido-5-pyridinio, 2,3-dichloro-1-oxido-5-pyridinio or 2-chlorothiazol-5-yl group,

A is especially a pyrid-3-yl, 2-chloropyrid-5-yl, 2-chloro-1-oxido-5-pyridinio or 2-chlorothiazol-5-yl group,

in particular A is a 2-chloropyrid-5-yl or, preferably, 2-chlorothiazol-5-yl group

(7) a compound of the formula (A), in which X is N-NO<sub>2</sub>;

(8) a compound of the formula (A), in which A is a 2-chlorothiazol-5-yl or 2-chloropyrid-5-yl group, R is C<sub>1</sub>-C<sub>4</sub>alkyl and X is N-NO<sub>2</sub>.

Compounds of the formula (A) which are named as preferred in the context of the invention are

- (A.1) (2-chloropyrid-5-ylmethyl)-3-methyl-4-nitroimino-perhydro-1,3,5-oxadiazine;
- (A.2) (2-chlorothiazol-5-ylmethyl)-3-ethyl-4-nitroimino-perhydro-1,3,5-oxadiazine;
- (A.3) 3-methyl-4-nitroimino-5-(1-oxido-3-pyridiniomethyl)-perhydro-1,3,5-oxadiazine;
- (A.4) (2-chloro-1-oxido-5-pyridiniomethyl)-3-methyl-4-nitroimino-perhydro-1,3,5-oxadiazine;
- (A.5) (2-chlorothiazol-5-ylmethyl)-3-methyl-4-nitroimino-perhydro-1,3,5-oxadiazine;
- (A.6) 3-methyl-5-(2-methylpyrid-5-ylmethyl)-4-nitroimino-perhydro-1,3,5-oxadiazine;
- (A.7) (2-chloropyrid-5-ylmethyl)-4-nitroimino-perhydro-1,3,5-oxadiazine;
- (A.8) (2-chlorothiazol-5-ylmethyl)-4-nitroimino-perhydro-1,3,5-oxadiazine, or

(A.9) (2-chloropyrid-5-ylmethyl)-3-ethyl-4-nitroimino-perhydro-1,3,5-oxadiazine.

5-(2-chlorothiazol-5-ylmethyl)-3-methyl-4-nitroimino-perhydro-1,3,5-oxadiazine is especially preferred in the context of the invention as the mixing partner of the formula (A). A composition which, in addition to the compound of the formula (A), also comprises only one further compound (I) to (CLXXXIV) having a pesticidal action is also preferred. Compositions which comprise pirimicarb in addition to a compound of the formula (A) are furthermore preferred. Compositions which comprise pyriproxyfen, fipronil, endosulfan or buprofezin in addition to a compound of the formula (A) are likewise preferred.

The active compound combination according to the invention preferably comprises the active compound of the formula (A) and one of the active compounds (I) to (CLXXXIV) in a mixing ratio of 100:1 to 1:6000, in particular 1:50 to 50:1, in particular in a ratio of between 1:20 and 20:1, in particular between 10:1 and 1:10, especially between 5:1 and 1:5, particularly preferably between 2:1 and 1:2, and also preferably between 4:1 and 2:1, especially in the ratio of 1:1, or 5:1, or 5:2, or 5:3, or 5:4, or 4:1, or 4:2, or 4:3, or 3:1, or 3:2, or 2:1, or 1:5, or 2:5, or 3:5, or 4:5, or 1:4, or 2:4, or 3:4, or 1:3, or 2:3, or 1:2, or, 1:600, or 1:300, or 1:150, or 1:35, or 2:35, or 4:35, or 1:75, or 2:75, or 4:75, or 1:6000, or 1:3000, or 1:1500, or 1:350, or 2:350, or 4:350, or 1:750, or 2:750, or 4:750. These ratios are understood as meaning weight ratios on the one hand, and also molar ratios on the other hand.

It has now been found, surprisingly, that the combination of the active compound of the formula (A) or of one of its salts with one of the active compounds (I) to (CLXXXIV) not only causes an additive increase in the action spectrum on the pests to be controlled, which is to be expected in principle, but that it achieves a synergistic effect which extends the action limits of both preparations under two aspects:

On the one hand, the rates of application of the compound of the formula (A) and of the individual compounds (I) to (CLXXXIV) are lowered for the same good action. On the other hand, the combined mixture also still has a high degree of pest control where the two individual substances have become completely inactive in the range of excessively low rates of application. This allows on the one hand a considerable extension to the spectrum of pests which can be controlled and on the other hand an increase in application reliability.

In addition to the actual synergistic action in respect of pesticidal activity, however, the compositions according to the invention also additionally have further surprising advantages which can likewise be described as synergistic in a broadened sense: thus, for example, it is possible to control pests which cannot be controlled or sufficiently effectively controlled with the individual compounds (A) or (I) to (CLXXXIV), and the compositions according to the invention show a better toleration by plants, i.e., for example, a reduced phytotoxicity, than the individual compounds (A) and (I) to (CLXXXIV). The insects can furthermore be controlled in their various stages of development, which sometimes is not the case with the individual compounds (A) and (I) to (CLXXXIV), since these compounds can be used, for example, only as adulticides or only as larvicides against quite specific larval stages. In addition, combinations of the compound (A) with certain compounds (I) to (CLXXXIV) show more favourable properties during grinding, mixing, storage and also spraying.

The compositions according to the invention are valuable preventively and/or curatively even at low rates of concentration in the field of pest control, while being tolerated by warm-blooded animals, fishes and plants, and have a very favourable biocidal spectrum. The compositions according to the invention are active against all or individual stages of development of normally sensitive and also resistant animal pests, such as insects and representatives of the order Acarina. The insecticidal and/or acaricidal action of the compositions according to the invention can manifest itself here directly, i.e. in a destruction of the pests, which occurs immediately or only after some time, for example during moulting, or indirectly, for example in a reduced oviposition and/or hatching rate, the good action corresponding to a rate of destruction (mortality) of at least 50 to 60%.

The animal pests include, for example:

from the order Lepidoptera

*Accleris* spp., *Adoxophyes* spp., *Aegeria* spp., *Agrotis* spp., *Alabama argillaceae*, *Amylois* spp., *Anticarsia gemmatilis*, *Archips* spp., *Argyrotaenia* spp., *Autographa* spp., *Busseola fusca*, *Cadra cautella*, *Carposina nipponensis*, *Chilo* spp., *Choristoneura* spp., *Clysia ambiguella*, *Cnaphalocrocis* spp., *Cnephasia* spp., *Cochylis* spp., *Coleophora* spp., *Crocidolomia binotalis*, *Cryptophlebia leucotreta*, *Cydia* spp., *Diatraea* spp., *Diparopsis castanea*, *Earias* spp., *Ephestia* spp., *Eucosma* spp., *Eupoecilia ambiguella*, *Euproctis* spp., *Euxoa* spp., *Grapholita* spp., *Hedya nubiferana*, *Heliothis* spp., *Hellula* spp., *Hyphantria cunea*, *Keiferia*

from the order Coleoptera, for example

from the order Orthoptera, for example

from the order Isoptera, for example

from the order Psocoptera, for example

**Liposcelis spp.;**

from the order Anoplura, for example

Haematopinus spp., Linognathus spp., Pediculus spp., Pemphigus spp. and Phylloxera spp.;

from the order Mallophaga, for example

**Damalinea spp. and Trichodectes spp.:**

from the order Thysanoptera, for example

Frankliniella spp., Hercinothrips spp., Taeniothrips spp., Thrips palmi, Thrips tabaci and Scirtothrips aurantii:

from the order Heteroptera, for example

Cimex spp., Distantiella theobroma, Dysdercus spp., Euchistus spp. Eurygaster spp. Leptocorisa spp., Nezara spp., Piesma spp., Rhodnius spp., Sahlbergella singularis, Scotinophara spp. and Triatoma spp.:

from the order Homoptera, for example



Aleurothrixus floccosus, Aleyrodes brassicae, Aonidiella spp., Aphididae, Aphis spp., Aspidiotus spp., Bemisia tabaci, Ceroplaster spp., Chrysomphalus aonidium, Chrysomphalus dictyospermi, Coccus hesperidum, Empoasca spp., Eriosoma lanigerum, Erythroneura spp., Gascardia spp., Laodelphax spp., Lecanium corni, Lepidosaphes spp., Macrosiphus spp., Myzus spp., Nephotettix spp., Nilaparvata spp., Paratoria spp., Pemphigus spp., Planococcus spp., Pseudaulacaspis spp., Pseudococcus spp., Psylla spp., Pulvinaria aethiopica, Quadraspidiotus spp., Rhopalosiphum spp., Saissetia spp., Scaphoideus spp., Schizaphis spp., Sitobion spp., Trialeurodes vaporariorum, Trioza erytrae and Unaspis citri;

from the order Hymenoptera, for example

Acromyrmex, Atta spp., Cephus spp., Diprion spp., Diprionidae, Gilpinia polytoma, Hoplocampa spp., Lasius spp., Monomorium pharaonis, Neodiprion spp., Solenopsis spp. and Vespa spp.;

from the order Diptera, for example

Aedes spp., Antherigona soccata, Bibio hortulanus, Calliphora erythrocephala, Ceratitis spp., Chrysomyia spp., Culex spp., Cuterebra spp., Dacus spp., Drosophila melanogaster, Fannia spp., Gastrophilus spp., Glossina spp., Hypoderma spp., Hyppobosca spp., Liriomyza spp., Lucilia spp., Melanagromyza spp., Musca spp., Oestrus spp., Orseolia spp., Oscinella frit, Pegomyia hyoscyami, Phorbia spp., Rhagoletis pomonella, Sciara spp., Stomoxys spp., Tabanus spp., Tannia spp. and Tipula spp.;

from the order Siphonaptera, for example

Ceratophyllus spp. and Xenopsylla cheopis;

from the order Thysanura, for example

Lepisma saccharina and

from the order Acarina, for example

Acarus siro, Aceria sheldoni, Aculus schlechtendali, Amblyomma spp., Argas spp., Boophilus spp., Brevipalpus spp., Bryobia praetiosa, Calipitimerus spp., Chorioptes spp., Dermanyssus gallinae, Eotetranychus carpini, Eriophyes spp., Hyalomma spp., Ixodes spp., Olygonychus pratensis, Ornithodoros spp., Panonychus spp., Phyllocoptruta oleivora, Polyphagotarsonemus latus, Psoroptes spp., Rhipicephalus spp., Rhizoglyphus spp., Sarcoptes spp., Tarsonemus spp. and Tetranychus spp..

Using the active compound mixtures according to the invention, in particular pests of the type mentioned which occur on plants, especially useful plants and ornamentals in agricultu-

re, in horticulture and in forestry, or on parts, such as fruit, blossom, foliage, stems, tubers, or roots, of such plants can be controlled, i.e. checked or destroyed, and in some cases parts of plants which additionally grow later are also still protected from these pests.

The pesticidal mixture according to the invention can advantageously be employed for pest control in cereals, such as maize or sorghum; in fruit, for example pome, stone and soft fruit, such as apples, pears, plums, peaches, almonds, cherries or berries, for example strawberries, raspberries and blackberries; in leguminous plants, such as beans, lentils, peas or soya; in oil-bearing crops, such as oilseed rape, mustard, poppy seed, olives, sunflower, coconut, castor, cacao or groundnuts; in gourds, such as pumpkins, cucumbers or melons; in fibre plants, such as cotton, flax, hemp or jute; in citrus fruits, such as oranges, lemons, grapefruits or mandarins; in vegetables, such as spinach, lettuce, asparagus, cabbage varieties, carrots, onions, tomatoes, potatoes or capsicum; in lauraceous plants, such as avocado, cinnamonium, camphor; or in tobacco, nuts, coffee, aubergines, sugar cane, tea, pepper, vines, hops, musaceous plants, natural rubber plants or ornamentals, in particular in maize, sorghum, pome and stone fruits, leguminous plants, cucurbits, cotton, citrus fruits, vegetables, aubergines, vines, hops or ornamentals, especially in maize, sorghum, apples, pears, plums, peaches, beans, peas, soya, olives, sunflower, coconut, cacao, groundnuts, cucumbers, pumpkins, citrus fruits, cabbage varieties, tomatoes, potatoes, vines or cotton, particularly preferably in vines, citrus fruits, apples, pears, tomatoes and cotton.

Other fields of use of the active compound mixtures according to the invention are protection of stored products and stocks and of material, and in the hygiene sector, in particular the protection of domestic animals and productive livestock against pests of the type mentioned.

Depending on the aims to be achieved and the given circumstances, the pesticides according to the invention are emulsifiable concentrates, suspension concentrates, directly sprayable or dilutable solutions, spreadable pastes, dilute emulsions, sprayable powders, soluble powders, dispersible powders, wettable powders, dusts, granules or encapsulations in polymeric substances which comprise the compound of the formula (A) or one of its salts and one of the other active compounds (I) to (CLXXXIV) according to the invention.

The active compounds are employed in these compositions in a pure form, the solid active compounds, for example, being employed in a specific particle size, or, preferably, together with - at least - one of the auxiliaries conventionally used in the art of formulation, such as extenders, for example solvents or solid carriers, or such as surface-active compounds (surfactants).

Solvents are, for example: non-hydrogenated or partly hydrogenated aromatic hydrocarbons, preferably fractions C<sub>8</sub> to C<sub>12</sub> of alkylbenzenes, such as xylene mixtures, alkylated naphthalenes or tetrahydronaphthalene, aliphatic or cycloaliphatic hydrocarbons, such as paraffins or cyclohexane, alcohols, such as ethanol, propanol or butanol, glycols and ethers and esters thereof, such as propylene glycol, dipropylene glycol ether, ethylene glycol or ethylene glycol monomethyl or -ethyl ether, ketones, such as cyclohexanone, isophorone or diacetone alcohol, strongly polar solvents, such as N-methylpyrrolid-2-one, dimethyl sulfoxide or N,N-dimethylformamide, water, non-epoxidized or epoxidized plant oils, such as non-epoxidized or epoxidized rapeseed, castor, coconut or soya oil, and silicone oils.

Solid carriers, for example for dusts and dispersible powders, which are as a rule used are natural rock powders, such as calcite, talc, kaolin, montmorillonite or attapulgite. Highly disperse silicic acids or highly disperse absorbent polymers can also be added to improve the physical properties. Granular, adsorptive granule carriers are porous types, such as pumice, crushed brick, sepiolite or bentonite, and non-adsorptive carrier materials are calcite or sand. A large number of granulated materials of inorganic or organic nature, in particular dolomite or comminuted plant residues, can furthermore be used.

Surface-active compounds are, depending on the nature of the active compound to be formulated, nonionic, cationic and/or anionic surfactants or surfactant mixtures having good emulsifying, dispersing and wetting properties. The surfactants listed below are to be regarded only as examples here; many other surfactants which are conventionally used in the art of formulation and are suitable according to the invention are described in the relevant literature.

Nonionic surfactants are, in particular, polyglycol ether derivatives of aliphatic or cycloaliphatic alcohols, saturated or unsaturated fatty acids and alkylphenols, which can contain 3 to 30 glycol ether groups and 8 to 20 carbon atoms in the (aliphatic) hydrocarbon radical and 6 to 18 carbon atoms in the alkyl radical of the alkylphenols. Water-soluble adducts,

containing 20 to 250 ethylene glycol ether groups and 10 to 100 propylene glycol ether groups, of polyethylene oxide on polypropylene glycol, ethylenediaminopolypropylene glycol and alkylpolypropylene glycol having 1 to 10 carbon atoms in the alkyl chain are furthermore suitable. The compounds mentioned usually comprise 1 to 5 ethylene glycol units per propylene glycol unit. Examples are nonylphenol-polyethoxyethanols, castor oil polyglycol ether, polypropylene/polyethylene oxide adducts, tributylphenoxypolyethoxyethanol, polyethylene glycol and octylphenoxypolyethoxyethanol. Fatty acid esters of polyoxyethylene-sorbitan, such as polyoxyethylene-sorbitan trioleate, are furthermore possible.

Cationic surfactants are, in particular, quaternary ammonium salts which contain, as substituents, at least one alkyl radical having 8 to 22 C atoms and, as further substituents, lower, non-halogenated or halogenated alkyl, benzyl or lower hydroxy alkyl radicals. The salts are preferably in the form of halides, methyl-sulfates or ethyl-sulfates. Examples are stearyltrimethylammonium chloride and benzyldi(2-chloroethyl)ethylammonium bromide.

Suitable inorganic surfactants can be both water-soluble soaps and water-soluble synthetic surface-active compounds. Suitable soaps are the alkali metal, alkaline earth metal and substituted and unsubstituted ammonium salts of higher fatty acids ( $C_{10}$ - $C_{22}$ ), such as the sodium or potassium salts of oleic or stearic acid, or of naturally occurring fatty acid mixtures, which can be obtained, for example, from coconut oil or tall oil; and furthermore also the fatty acid methyl-aurine salts. However, synthetic surfactants are more frequently used, in particular fatty sulfonates, fatty sulfates, sulfonated benzimidazole derivatives or alkylarylsulfonates. The fatty sulfonates and sulfates are as a rule in the form of alkali metal, alkaline earth metal or substituted or unsubstituted ammonium salts, and in general contain an alkyl radical having 8 to 22 C atoms, alkyl also including the alkyl moiety of acyl radicals; examples are the sodium or calcium salt of ligninsulfonic acid, of dodecylsulfuric acid ester or of a fatty alcohol sulfate mixture prepared from naturally occurring fatty acids. These also include the salts of sulfuric acid esters and sulfonic acids of fatty alcohol-ethylene oxide adducts. The sulfonated benzimidazole derivatives preferably contain two sulfonic acid groups and one fatty acid radical having about 8 to 22 C atoms. Alkylarylsulfonates are, for example, the sodium, calcium or triethanolammonium salts of dodecylbenzenesulfonic acid, of dibutyl-naphthalenesulfonic acid or of a naphthalenesulfonic acid-formaldehyde condensati-

on product. Corresponding phosphates, such as salts of the phosphoric acid ester of a p-nonylphenol-(4-14)-ethylene oxide adduct or phospholipids, are furthermore also suitable.

The compositions as a rule comprise 0.1 to 99%, in particular 0.1 to 95%, of a mixture of the active compound of the formula (A) with one of the active compounds (I) to (CLXXXIV), and 1 to 99.9%, in particular 5 to 99.9%, of - at least - one solid or liquid auxiliary, where as a rule 0 to 25%, in particular 0.1 to 20%, of the compositions can be surfactants (% is in each case per cent by weight). While concentrated compositions are rather preferred as commercial goods, the end user as a rule uses dilute compositions which have considerably lower active compound concentrations. Preferred compositions have, in particular, the following composition (% = per cent by weight):

Emulsifiable concentrates:

Active compound mixture	1 to 90%, preferably 5 to 20%
Surfactant:	1 to 30%, preferably 10 to 20 %
Solvent:	5 to 98%, preferably 70 to 85%

Dusts:

Active compound mixture:	0.1 to 10%, preferably 0.1 to 1%
Solid carrier:	99.9 to 90%, preferably 99.9 to 99%

Suspension concentrates:

Active compound mixture:	5 to 75%, preferably 10 to 50%
Water:	94 to 24%, preferably 88 to 30%
Surfactant:	1 to 40%, preferably 2 to 30%

Wettable powders:

Active compound mixture:	0.5 to 90%, preferably 1 to 80%
Surfactant:	0.5 to 20%, preferably 1 to 15%
Solid carrier:	5 to 99%, preferably 15 to 98%

Granules:

Active compound mixture:	0.5 to 30%, preferably 3 to 15%
Solid carrier:	99.5 to 70%, preferably 97 to 85%

The compositions according to the invention can also comprise other solid or liquid auxiliaries, such as, stabilizers, for example non-epoxidized or epoxidized plant oils (for example epoxidized coconut oil, rape oil or soya oil), defoamers, for example silicone oil, preservatives, viscosity regulators, binders and/or tackifiers, as well as fertilizers or other active compounds for achieving specific effects, for example bactericides, fungicides, nematocides, molluscicides or herbicides.

The compositions according to the invention are prepared in a known manner, in the absence of auxiliaries, for example, by grinding, sieving and/or pressing a solid active compound or active compound mixture, for example to a particular particle size, and in the presence of at least one auxiliary, for example, by intimate mixing and/or grinding of the active compound or active compound mixture with the auxiliary or auxiliaries. The invention therefore also relates to the process for the preparation of the compositions.

Mixtures of a compound of the formula (A) with one or more of the compounds (I) to (CLXXXIV) are preferably employed with the auxiliaries conventionally used in the art of formulation, and are therefore processed, for example, to emulsifiable concentrates, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts and granules, and also to encapsulations in, for example, polymeric substances, in a known manner. The methods of application, such as spraying, atomizing, dusting, wetting, scattering or pouring are chosen, as is the nature of the composition, according to the aims to be achieved and the given circumstances.

The invention furthermore relates to the methods of application for the compositions, i.e. the methods for controlling pests of the type mentioned, such as spraying, atomizing, dusting, spreading, dressing, scattering or pouring, to be chosen according to the aims to be achieved and the given circumstances, and the use of the compositions for controlling pests of the type mentioned. Typical rates of concentration here are between 0.1 and 1000 ppm, preferably between 0.1 and 500 ppm, of active compound. The rate of application can vary within wide limits and depends on the nature of the soil, the nature of the application, (leaf application; seed dressing, application in the seed furrow), the crop plant, the pest to be controlled, the particular climatic circumstances which prevail and other factors determined by the nature of the application, the time of application and the target crop. The rates of

application per hectare are in general 1 to 2000 g of active compound per hectare, in particular 10 to 1000 g/ha, preferably 20 to 600 g/ha.

A preferred method of application in the field of crop protection is application to the foliage of the plants (leaf application), where the frequency of application and rate of application can be chosen according to the risk of infestation by the particular pest. However, the active compound can also enter the plants via the root system (systemic action) by impregnating the locus of the plants with a liquid composition or introducing the active compounds in solid form into the locus of the plants, for example into the soil, for example in the form of granules (soil application). In paddy rice crops, such granules can be metered onto the flooded rice field.

The compositions according to the invention are also suitable for the protection of plant propagation material, for example seeds, such as fruit, tubers, or grain, or plant seedlings, against animal pests. The propagation material can be treated with the composition before being brought out, for example seed can be dressed before sowing. The active compounds according to the invention can also be applied to the seed grains (coating) either by impregnating the grains in a liquid composition or coating them with a solid composition. The composition can also be applied to the site where the propagation material is brought out, during broadcasting, for example into the seed furrow during sowing. The invention furthermore relates to the treatment methods for plant propagation material and the plant propagation material treated in this way.

The following examples serve to illustrate the invention. They do not limit the invention.

Formulation examples (% = % by weight, active compound ratios = weight ratios)

Example F1: Emulsion concentrates

	a)	b)	c)
Active compound mixture (Compound (A) :			
Compound (I) to (CLXXXIV) = 1 : 1)	25 %	40 %	50 %
Ca dodecylbenzenesulfonate	5 %	8 %	6 %
Castor oil polyethylene glycol ether (36 mol of EO)	5 %		
Tributylphenolpolyethylene glycol ether (30 mol of EO)		12 %	4 %
Cyclohexanone		15 %	20 %
Xylene mixture	65 %	25 %	20 %

EO is the degree of ethoxylation of castor oil or tributylphenol.

Emulsions of any desired concentration can be prepared from such concentrates by dilution with water.

Example F2: Solutions

	a)	b)	c)	d)
Active compound mixture (Compound (A) :				
Compound (I) to (CLXXXIV) = 1 : 3)	80 %	10 %	5 %	95 %
Ethylene glycol monomethyl ether		20 %		
Polyethylene glycol molecular wt. 400	-	70 %		
N-methyl-2-pyrrolidone	20 %	-		
Epoxidized coconut oil			1 %	5 %
Benzine (boiling limits 160-190°C)			94 %	

The solutions are suitable for use in the form of tiny drops.

Example F3: Granules

	a)	b)	c)	d)
Active compound mixture (Compound (A) :				
Compound (I) to (CLXXXIV) = 3 : 1)	5 %	10 %	8 %	21 %
Kaolin	94 %		79 %	54 %
Highly disperse silicic acid	1 %		13 %	7 %
Attapulgate		90 %		18 %

The active compounds are dissolved together in methylene chloride, the solution is sprayed onto the carrier and the solvent is then evaporated off in vacuo.

Example F4: Dusts

	a)	b)
Active compound mixture (1 : 1)	2 %	5 %
Highly disperse silicic acid	1 %	5 %
Talc	97 %	
Kaolin		90 %

Intimate mixing of the carriers with the active compounds gives ready-to-use dusts.

05716655-11000



Example F5: Wettable powders

	a	b)	c)
Active compound mixture (Compound (A) :			
Compound (I) to (CLXXXIV) = 4 : 1)	25 %	50 %	75 %
Sodium ligninsulfonate	5 %	5 %	
Sodium lauryl sulfate	3 %		5 %
Sodium diisobutylphenylsulfonate		6 %	10 %
Octylphenol polyethylene glycol-ether (7-8 mol of EO)		2 %	-
Highly disperse silicic acid	5 %	10 %	10 %
Kaolin	62 %	27 %	-

The active compounds are mixed with the additives and ground thoroughly in a suitable mill. Wettable powders which can be diluted with water to give suspensions of any desired concentration are obtained.

Example F6: Emulsion concentrate

Active compound mixture (1 : 2)	10 %
Octylphenol polyethylene glycol ether	
(4-5 mol of EO)	3 %
Calcium dodecylbenzenesulfonate	3 %
Castor oil polyglycol ether	
(36 mol of EO)	4 %
Cyclohexanone	30 %
Xylene mixture	50 %

Emulsions of any desired concentration can be prepared from this concentrate by dilution with water.

Example F7: Dusts

	a)	b)
Active compound mixture (Compound (A) :		
Compound (I) to (CLXXXIV) = 2 : 5)	5%	8 %
Talc	95 %	-
Kaolin		92 %

Ready-to-use dusts are obtained by mixing the active compounds with the carrier and grinding the mixture on a suitable mill.

Example F8: Extruded granules

Active compound mixture (2 : 1)	10 %
Sodium ligninsulfonate	2 %
Carboxymethylcellulose	1 %
Kaolin	87 %

The active compounds are mixed with the additives and the mixture is ground and moistened with water. This mixture is extruded, granulated and then dried in a stream of air.

Example F9: Coated granules

Active compound mixture (1 : 1)	3 %
Polyethylene glycol (molecular wt. 200)	3 %
Kaolin	94 %

The finely ground active compounds are applied uniformly to the kaolin, which is moistened with polyethylene glycol, in a mixer. Dust-free coated granules are obtained in this way.

Example F10: Suspension concentrate

Active compound mixture (1 : 1)	40 %
Ethylene glycol	10 %
Nonylphenol polyethylene glycol ether (15 mol of EO)	6 %
Sodium ligninsulfonate	10 %
Carboxymethylcellulose	1 %
37% aqueous formaldehyde solution	0,2 %
Silicone oil in the form of a 75% aqueous emulsion	0.8 %
Water	32 %

The finely ground active compounds are mixed intimately with the additives. A suspension concentrate from which suspensions of any desired concentration can be prepared by dilution with water is thus obtained.

It is often more practical to formulate the active compound of the formula (A) and one of the mixing partners (I) to (CLXXXIV) individually and then to bring them together in the applicator in the desired mixing ratio as "tank mixing" in the water only shortly before application.

Biological Examples (% = per cent by weight, unless stated otherwise)

A synergistic effect is always present if the action EA of the combination of an active compound of the formula (A) with one of the active compounds (I) to (CLXXXIV) is greater than the sum of the action of the active compounds applied individually:

$$EA_1 > X + Y \quad (B)$$

However, the pesticidal action EA to be expected for a given combination of two pesticides can also be calculated as follows (cf. COLBY, S.R., "Calculating synergistic and antagonistic response of herbicide combinations", Weeds 15, Pages 20-22, 1967).

In this equation:

$$EA_2 = X + \frac{Y (100 - X)}{100}$$

X = Per cent mortality on treatment with the compound of the formula (A) at a rate of application of p kg per hectare compared with the untreated control (= 0 %).

Y = Per cent mortality on treatment with a compound (I) to (CLXXXIV) at a rate of application of q kg per hectare compared with the untreated control.

EA = Expected pesticidal action (per cent mortality compared with the untreated control) after treatment with the compound of the formula (A) and a compound (I) to (CLXXXIV) at a rate of application of p + q kg of active compound per hectare.

If the action actually observed is greater than the expected value EA, synergism exists.

Example B1: Action against Bemisia tabaci

Dwarf bean plants are placed in gauze cages and populated with adults of Bemisia tabaci. After oviposition has taken place, all the adults are removed. 10 days later, the plants with the nymphs on them are sprayed with an aqueous suspension spray liquor comprising 50 ppm of the active compound mixture. After a further 14 days, the percentage hatching of the eggs is evaluated in comparison with untreated control batches.

In this test, the combinations of an active compound of the formula (A) with one of the active compounds (I) to (CLXXXIV) show a synergistic effect. In particular, a suspension spray liquor which comprises 40 ppm of the compound (A.5) and 10 ppm of the compound (II) shows an activity of over 80%.

Example B2: Action against *Spodoptera littoralis* caterpillars

Young soya plants are sprayed with an aqueous emulsion spray liquor comprising 360 ppm of the active compound mixture. After the spray coating has dried on, the soya plants are populated with 10 caterpillars of the third stage of *Spodoptera littoralis* and placed in a plastic container. Evaluation takes place 3 days later. The percentage reduction in the population and the percentage reduction in feeding damage (% action) are determined from comparison of the number of dead caterpillars and of the feeding damage on the treated plants to those on the untreated plants.

In this test, combinations of an active compound of the formula (A) with one of the active compounds (I) to (CLXXXIV) show a synergistic effect. In particular, a suspension spray liquor which comprises 200 ppm of the compound (A.2) and 160 ppm of the compound (XXIX) and a suspension spray liquor which comprises 120 ppm of the compound (A) and 240 ppm of the compound (XXIV) show an activity of over 80%.

Example B3: Ovizidal action on *Lobesia botrana*

*Lobesia botrana* eggs deposited on filter paper are immersed for a short time in an acetone-aqueous test solution comprising 400 ppm of the active compound mixture to be tested. After the test solution has dried on, the eggs are incubated in Petri dishes. After 6 days, the percentage hatching of the eggs is evaluated in comparison with untreated control batches (% reduction in hatching).

In this test, combinations of an active compound of the formula (A) with one of the active compounds (I) to (CLXXXIV) show a synergistic effect. In particular, a suspension spray liquor which comprises 300 ppm of the compound (A.5) and 100 ppm of the compound (III) and a suspension spray liquor which comprise 200 ppm of the compound (A.2) and 200 ppm of the compound (LVII) show an activity of over 80%.

Example B4: Ovizidal action on *Heliothis virescens*

*Heliothis virescens* eggs deposited on filter paper are immersed for a short time in an acetone-aqueous test solution comprising 400 ppm of the active compound mixture to be tested. After the test solution has dried on, the eggs are incubated in Petri dishes. After 6 days, the percentage hatching of the eggs is evaluated in comparison with untreated control batches (% reduction in hatching).

In this test, combinations of an active compound of the formula (A) with one of the active compounds (I) to (CLXXXIV) show a synergistic effect. In particular, a suspension spray liquor which comprises 240 ppm of the compound (A.1) and 160 ppm of the compound (I) and a suspension spray liquor which comprise 100 ppm of the compound (A.5) and 300 ppm of the compound (I) show an activity of over 80%.

Example B5: Action against *Plutella xylostella* caterpillars

Young cabbage plants are sprayed with an aqueous emulsion spray liquor which comprises 440 ppm of the active compound. After the spray coating has dried on, the cabbage plants are populated with ten caterpillars of the third stage of *Plutella xylostella* and placed in a plastic container. The evaluation takes place 3 days later. The percentage reduction in population and percentage reduction in feeding damage (% action) are determined from the comparison of the number of dead caterpillars and of the feeding damage on the treated plants to those on the untreated plants.

In this test, combinations of an active compound of the formula (A) with one of the active compounds (I) to (CLXXXIV) show a synergistic effect. In particular, a suspension spray liquor which comprises 40 ppm of the compound (A.5) and 400 ppm of the compound (XI) and a suspension spray liquor which comprises 220 ppm of the compound (A.7) and 220 ppm of the compound (L) show an activity of over 80%.


Example B6: Action against *Myzus persicae*

Pea seedlings are infected with *Myzus persicae*, subsequently sprayed with a spray liquor comprising 400 ppm of the active compound and incubated at 20°C. An evaluation takes place 3 and 6 days later. The percentage reduction in the population (% action) is determined from the comparison of the number of dead aphids on the treated plants to that on the untreated plants.

In this test, combinations of an active compound of the formula (A) with one of the active compounds (I) to (CLXXXIV) show a synergistic effect. In particular, a suspension spray liquor which comprises 1 ppm of the compound (A.5) and 40 ppm of pirimicarb and a suspension spray liquor which comprises 0.2 ppm of the compound (A.5) and 75 ppm of pirimicarb show an activity of over 80%\*.

WHAT IS CLAIMED IS:

1. A composition for controlling insects or representatives of the order Acarina, which comprises a combination of variable amounts of one or more compounds of the formula



in which

A is an unsubstituted or, depending on the possibility of substitution on the ring system, mono- to tetrasubstituted, aromatic or non-aromatic monocyclic or bicyclic heterocyclic radical, in which the substituents of A are chosen from the group consisting of C<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>3</sub>alkoxy, halogen, halo-C<sub>1</sub>-C<sub>3</sub>alkyl, cyclopropyl, halocyclopropyl, C<sub>2</sub>-C<sub>3</sub>alkenyl, C<sub>2</sub>-C<sub>3</sub>alkynyl, halo-C<sub>2</sub>-C<sub>3</sub>alkenyl, halo-C<sub>2</sub>-C<sub>3</sub>alkynyl, halo-C<sub>1</sub>-C<sub>3</sub>alkoxy, C<sub>1</sub>-C<sub>3</sub>alkylthio, Halo-C<sub>1</sub>-C<sub>3</sub>alkylthio, allyloxy, propargyloxy, allylthio, propargylthio, haloallyloxy, haloallylthio, cyano and nitro;

R is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl or C<sub>2</sub>-C<sub>6</sub>alkynyl; and

X is N-NO<sub>2</sub> or N-CN,

in the free form or in salt form, if appropriate tautomers, in the free form or salt form, and one or more of the compounds:

(I) aldicarb;	(XIV) zeta-Cypermethrin;	(XXVII) methamidophos;
(II) azinphos-methyl;	(XV) deltamethrin;	(XXVIII) methomyl;
(III) benfuracarb;	(XVI) diflubenzuron;	(XXIX) mevinphos;
(IV) bifenthrin;	(XVII) endosulfan;	(XXX) parathion;
(V) buprofezin;	(XVIII) ethiofencarb;	(XXXI) parathion-methyl;
(VI) carbofuran;	(XIX) fenitrothion;	(XXXII) phosalone;
(VII) dibutylaminothio;	(XX) fenobucarb;	(XXXIII) pirimicarb;
(VIII) cartap;	(XXI) fenvalerate;	(XXXIV) propoxur;
(IX) chlorofluazuron;	(XXII) formothion;	(XXXV) teflubenzuron;
(X) chloropyrifos;	(XXIII) methiocarb;	(XXXVI) terbufos;
(XI) cyfluthrin;	(XXIV) heptenophos;	(XXXVII) triazamate;
(XII) lambda-cy-halothrin;	(XXV) imidacloprid;	(XXXVIII) abamectin;
(XIII) alpha-cypermethrin;	(XXVI) isoprocarb;	(XXXIX) fenobucarb;

- |  |                           |                         |
|--|---------------------------|-------------------------|
| (XL) tebufenozide;   | (XLIV) fenpyroximate;     | (XLVIII) pyrimidifen;   |
| (XLI) fipronil;  | (XLV) pyridaben;          | (XLIX) nitenpyram;      |
| (XLII) beta-cyfluthrin;  | (XLVI) fenazaquin;        | (L) NI-25, acetamiprid; |
| (XLIII) silafluofen;   | (XLVII) pyriproxyfen;     |                         |
| (LI) avermectin B <sub>1</sub> (abamectin);                                      |                           |                         |
| (LII) a plant extract which is active against insects;                           |                           |                         |
| (LIII) a preparation which comprises nematodes which are active against insects; |                           |                         |
| (LIV) a preparation obtainable from <i>Bacillus subtilis</i> ;                   |                           |                         |
| (LV) a preparation which comprises fungi which are active against insects;       |                           |                         |
| (LVI) a preparation which comprises viruses which are active against insects;    |                           |                         |
| (LVII) AC 303 630;   | (LXXIX) chloethocarb;     | (CII) ethoprophos;      |
| (LVIII) acephate;  | (LXXX) chloroethoxyfos;   | (CIII) etrimphos;       |
| (LIX) acrinathrin;   | (LXXXI) chloromephos;     | (CIV) fenamiphos;       |
| (LX) alanycarb;  | (LXXXII) cis-res-methrin; | (CV) fenbutatin oxide;  |
| (LXI) alphamethrin;  | (LXXXIII) clocythrin;     | (CVI) fenothiocarb;     |
| (LXII) amitraz;  | (LXXXIV) clofentezin;     | (CVII) fenpropathrin;   |
| (LXIII) AZ 60541;  | (LXXXV) cyanophos;        | (CVIII) fenpyrad;       |
| (LXIV) azinphos A;   | (LXXXVI) cycloprothrin;   | (CIX) fenthion;         |
| (LXV) azinphos M;  | (LXXXVII) cyhexatin;      | (CX) fluazinam;         |
| (LXVI) azocyclotin;  | (LXXXVIII) demeton M;     | (CXI) flucycloxuron;    |
| (LXVII) bendiocarb;  | (LXXXIX) demeton S;       | (CXII) flucythrinate;   |
| (LXVIII) bensultap;  | (XC) demeton-S-methyl;    | (CXIII) flufenoxuron;   |
| (LXIX) betacyfluthrin;   | (XCI) dichlofenthion;     | (CXIV) flufenprox;      |
| (LXX) BPMC;  | (XCII) dicliphos;         | (CXV) fonophos;         |
| (LXXI) brofenprox;   | (XCIII) diethion;         | (CXVI) fosthiazate;     |
| (LXXII) bromophos A;   | (XCIV) dimethoate;        | (CXVII) fubfenprox;     |
| (LXXIII) bufencarb;  | (XCV) dimethylvinphos;    | (CXVIII) HCH;           |
| (LXXIV) butocarboxin;  | (XCVI) dioxathion;        | (CXIX) hexaflumuron;    |
| (LXXV) butylpyridaben;   | (XCVII) edifenphos;       | (CXX) hexythiazox;      |
| (LXXVI) cadusafos;   | (XCVIII) emamectin;       | (CXXI) iprobenfos;      |
| (LXXVII) carbaryl;   | (XCIX) esfenvalerate;     | (CXXII) isofenphos;     |
| (LXXVIII) carbopheno-  | (C) ethion;               | (CXXIII) isoxathion;    |
| thion;   | (CI) ethofenprox;         | (CXXIV) ivermectin;     |

(CXXV) lambda-cyhalothrin;	(CXLV) pirimiphos A;	(CLXVI) thiodicarb;
(CXXVI) malathion;	(CXLVI) promecarb;	(CLXVII) thiofanox;
(CXXVII) mecarbam;	(CXLVII) propaphos;	(CLXVIII) thionazin;
(CXXVIII) mesulfenphos;	(CXLVIII) prothiofos;	(CLXIX) thuringiensin;
(CXXIX) metaldehyde;	(CXLIX) prothoate;	(CLXX) tralomethrin;
(CXXX) metolcarb;	(CL) pyrachlophos;	(CLXXI) triarthen;
(CXXXI) milbemectin;	(CLI) pyrada-phenthion;	(CLXXII) triazophos;
(CXXXII) moxidectin;	(CLII) pyresmethrin;	(CLXXIII) triazuron;
(CXXXIII) naled;	(CLIII) pyrethrum;	(CLXXIV) trichlorofon;
(CXXXIV) NC 184;	(CLIV) RH 5992;	(CLXXV) triflumuron;
(CXXXV) omethoate;	(CLV) salithion;	(CLXXVI) trimethacarb;
(CXXXVI) oxamyl;	(CLVI) sebufos;	(CLXXVII) vamidothion;
(CXXXVII) oxydeme-	(CLVII) sulfotep;	(CLXXVIII) xylylcarb;
thion M;	(CLVIII) sulprofos;	(CLXXIX) YI 5301/5302;
(CXXXVIII) oxydeprofos;	(CLIX) tebufenpyrad;	(CLXXX) zetamethrin;
(CXXXIX) permethrin;	(CLX) tebupirimphos;	(CLXXXI) DPX-MP062;
(CXL) phenthoate;	(CLXI) tefluthrin;	(CLXXXII) RH-2485;
(CXLI) phorate;	(CLXII) temephos;	(CLXXXIII) D 2341; or
(CXLII) phosmet;	(CLXIII) terbam;	(CLXXXIV) XMC (3,5-
(CXLIII) phoxim;	(CLXIV) tetrachloro-	xylylmethylcarbamate),
(CXLIV) pirimiphos M;	vinphos;	
	(CLXV) thiafenox;	

and at least one auxiliary.

2. A composition according to claim 1, in which, in the compound of the formula (A), R is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl or C<sub>2</sub>-C<sub>6</sub>alkynyl.

3. A composition according to claim 1 or 2, in which, in the compound of the formula (A), the cyclic base skeleton of A contains 2 to 4 double bonds.

4. A composition according to any one of claims 1 to 3, in which, in the compound of the formula (A), the cyclic base skeleton of A contains 1 up to and including 4 heteroatoms.




5. A composition according to any one of claims 1 to 4, in which, in the compound of the formula (A), the cyclic base skeleton of A contains 1, 2 or 3 heteroatoms, chosen from the group consisting of oxygen, sulfur and nitrogen, not more than one of the heteroatoms contained in the cyclic base skeleton being an oxygen or a sulfur atom.
6. A composition according to any one of claims 1 to 5, in which, in the compound of the formula (A) the cyclic base skeleton of A is mono- or disubstituted by substituents chosen from the group consisting of halogen and C<sub>1</sub>-C<sub>3</sub>alkyl.
7. A composition according to any one of claims 1 to 6, in which, in the compound of the formula (A), the cyclic base skeleton of A is a pyridyl, 1-oxidopyridinio or thiazolyl group.
8. A composition according to any one of claims 1 to 7, in which, in the compound of the formula (A), X is N-NO<sub>2</sub>.
9. A composition according to claim 1, which comprises either
- (A.1) (2-chloropyrid-5-ylmethyl)-3-methyl-4-nitroimino-perhydro-1,3,5-oxadiazine;
  - (A.2) (2-chlorothiazol-5-ylmethyl)-3-ethyl-4-nitroimino-perhydro-1,3,5-oxadiazine;
  - (A.3) 3-methyl-4-nitroimino-5-(1-oxido-3-pyridiniomethyl)-perhydro-1,3,5-oxadiazine;
  - (A.4) (2-chloro-1-oxido-5-pyridiniomethyl)-3-methyl-4-nitroimino-perhydro-1,3,5-oxadiazine;
  - (A.5) (2-chlorothiazol-5-ylmethyl)-3-methyl-4-nitroimino-perhydro-1,3,5-oxadiazine;
  - (A.6) 3-methyl-5-(2-methylpyrid-5-ylmethyl)-4-nitroimino-perhydro-1,3,5-oxadiazine;
  - (A.7) (2-chloropyrid-5-ylmethyl)-4-nitroimino-perhydro-1,3,5-oxadiazine;
  - (A.8) (2-chlorothiazol-5-ylmethyl)-4-nitroimino-perhydro-1,3,5-oxadiazine; or
  - (A.9) (2-chloropyrid-5-ylmethyl)-3-ethyl-4-nitroimino-perhydro-1,3,5-oxadiazine.
10. A composition according to any one of claims 1 to 9, which comprises 5-(2-chlorothiazol-5-ylmethyl)-3-methyl-4-nitroimino-perhydro-1,3,5-oxadiazine.
11. A composition according to any one of claims 1 to 10, which comprises only one of the compounds (I) to (CLXXXIV).
12. A composition according to any one of claims 1 to 11 which comprises pyriproxyfen.
13. A composition according to any one of claims 1 to 11, which comprises fipronil.

14. A composition according to any one of claims 1 to 11, which comprises endosulfan.
15. A composition according to any one of claims 1 to 11, which comprises buprofezin.
16. A composition according to any one of claims 1 to 11, which comprises pirimicarb.
17. A method of controlling pests, which comprises applying a composition as defined in any one of claims 1 to 16, to the pests or their environment.
18. A method according to claim 17, for the protection of plant propagation material, which comprises treating the plant propagation material or the site where the propagation material is brought out.
19. A process for the preparation of a composition comprising at least one auxiliary as defined in any one of claims 1 to 16, which comprises intimately mixing the active compounds with the auxiliary or auxiliaries.
20. Plant propagation material treated by the method defined in claim 18.
21. The use of a composition as defined in any one of claims 1 to 16, in a method as defined in claim 17 or 18.
22. The use of a compound of the formula (A), in the free form or in an agrochemically usable salt form, for the preparation of a composition as defined in any one of claims 1 to 16.

## Pesticidal composition

### Abstract:

Composition for controlling insects and representatives of the order Acarina, which comprises a combination of variable amounts of one or more compounds of the formula



in which

A is an unsubstituted or, depending on the possibility of substitution on the ring system, mono- to tetrasubstituted, aromatic or non-aromatic monocyclic or bicyclic heterocyclic radical, in which the substituents of A can be chosen from the group consisting of C<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>3</sub>alkoxy, halogen, halo-C<sub>1</sub>-C<sub>3</sub>alkyl, cyclopropyl, halocyclopropyl, C<sub>2</sub>-C<sub>3</sub>alkenyl, C<sub>2</sub>-C<sub>3</sub>alkynyl, halo-C<sub>2</sub>-C<sub>3</sub>alkenyl, halo-C<sub>2</sub>-C<sub>3</sub>alkynyl, halo-C<sub>1</sub>-C<sub>3</sub>alkoxy, C<sub>1</sub>-C<sub>3</sub>alkylthio, Halo-C<sub>1</sub>-C<sub>3</sub>alkylthio, allyloxy, propargyloxy, allylthio, propargylthio, haloallyloxy, haloallylthio, cyano and nitro;

R is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl or C<sub>2</sub>-C<sub>6</sub>alkynyl; and

X is N-NO<sub>2</sub> or N-CN,

in the free form or in salt form, if appropriate tautomers, in the free form or salt form, and one or more of the compounds (I) to (CLXXXIV) mentioned according to the invention and at least one auxiliary.

A method of controlling pests, a process for the preparation of the composition, its use and plant propagation material treated with it, and the use of the compound of the formula (A) for the preparation of the composition, are described.

**DECLARATION AND POWER OF ATTORNEY FOR UNITED STATES PATENT APPLICATION**

☒ Original

☐ Supplemental

☐ Substitute

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name, and

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if more than one name is listed below) of the subject matter which is claimed and for which a United States patent is sought on the invention entitled

**PESTICIDAL COMPOSITIONS**

the specification of which:

☐ is attached hereto.

☐ was filed on \_\_\_\_\_ as Application No. \_\_\_\_\_  
(day/month/year)

and, if this box (☐) contains an ✕

☐ was amended on \_\_\_\_\_  
(day/month/year)

☒ was filed as Patent Cooperation Treaty international Application No.

PCT/EP 97/01926 on 17/04/1997  
(day/month/year)

and, if this box (☐) contains an ✕

☐ entered the national stage in the United States and was accorded Application No. \_\_\_\_\_

and, if this box (☐) contains an ✕

☐ was amended, subsequent to entry into the national stage, on \_\_\_\_\_  
(day/month/year)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above and, if this application was filed as a Patent Cooperation Treaty international application, by any amendments made during the international stage (including any made under Patent Cooperation Treaty Rule 91, Article 19 and Article 34).

I acknowledge my duty to disclose all information which is known by me to be material to the patentability of this application as defined in 37 C.F.R. § 1.56.

I hereby claim the benefit under 35 U.S.C. §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate listed below and under 35 U.S.C. §365(a) of any Patent Cooperation Treaty international application(s) designating at least one country other than the United States listed below and have also listed below any foreign application(s) for patent or inventor's certificate and Patent Cooperation Treaty international application(s) designating at least one country other than the United States for the same subject matter and having a filing date before that of the application the priority of which is claimed for that subject matter:

COUNTRY/REGION (OR P.C.T.)	APPLICATION No.	FILING DATE (day/month/year)	PRIORITY CLAIMED	
Switzerland	1082/96	29/04/96	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No

I hereby claim the benefit under 35 U.S.C. § 119 (e) of any United States provisional application(s) listed below:

APPLICATION NO.	FILING DATE (day/month/year)
-----------------	---------------------------------

I hereby claim the benefit under 35 U.S.C. §120 of any United States application(s) listed below and under 35 U.S.C. §365(c) of any Patent Cooperation Treaty international application(s) designating the United States listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in said prior application(s) in the manner required by the first paragraph of 35 U.S.C. §112, I acknowledge my duty to disclose all information known to me to be material to patentability as defined in 37 C.F.R. §1.56 which became available between the filing date(s) of the prior application(s) and the national or Patent Cooperation Treaty international filing date of this application:

United States Application No.	United States Filing Date (day/month/year)	Status (Pending, Abandoned or U.S. Patent No.)	International Application No.	and Filing Date
----------------------------------	--	--	-------------------------------------	--------------------

Full name of second  
joint inventor, if any

**Peter MAIENFISCH**

Inventor's signature



Date

14. 10. 98  
(day/month/year)

Residence

**4118 Rodersdorf, Switzerland**

Citizenship

**Swiss**


Post Office Address

**Aegertenstrasse 21  
4118 Rodersdorf  
Switzerland**

Full name of third  
joint inventor, if any

**Peter WYSS**

Inventor's signature



Date

19. 10. 98  
(day/month/year)

Residence

**2748 Souboz, Switzerland**

Citizenship

**Swiss**

Post Office Address

**Do Tia 14  
2748 Souboz  
Switzerland**